Supplementary Appendix

This appendix has been provided by the authors to give readers additional information about their work.

Supplement to: Nelson DA, Deuster PA, Carter R III, Hill OT, Wolcott VL, Kurina LM. Sickle cell trait, rhabdomyolysis, and mortality among U.S. army soldiers. N Engl J Med 2016;375:435-42. DOI: 10.1056/NEJMoa1516257

Supplementary Appendix

Table of Contents	Page
List of investigators	2
Table S1. Methods employed by the U.S. Army to prevent exertional	3
apse events, including heat injury and rhabdomyolysis, during	
training	
References	5

List of Investigators

D. Alan Nelson, Ph.D.; U.S. Army Office of the Surgeon General

Patricia A. Deuster, Ph.D.; Consortium for Health and Military Performance, A DoD Center of Excellence; Department of Military and Emergency Medicine; Uniformed Services University

Robert Carter III, Ph.D.; U.S. Army Research Institute of Surgical Research, Fort Sam Houston & Department of Emergency Medicine, University of Texas Health Science Center at San Antonio

O. Hill, Ph.D.; Extremity Trauma and Amputation Center of Excellence, Center for the Intrepid, Brooke Army Medical Center

Vickee L. Wolcott, Ph.D.; Army-Baylor University Graduate Program in Health and Business Administration

Lianne M. Kurina, Ph.D.; Department of Medicine, Division of Primary, Preventive, and Community Medicine, Stanford University School of Medicine; 450 Serra Mall, Bldg 20, Stanford, CA 94305-2160; (650) 725-7531; lkurina@stanford.edu (corresponding author)

Table S1. Methods employed by the US Army to prevent exertional collapse events during training, including heat injury and rhabdomyolysis.

Leader, clinician and individual prevention measures¹

Identify and assess hazards

- Check the recent and anticipated environmental conditions and associated heat stress levels, and determine unique group and individual risk factors
- Use the "HEAT" acronym:
 - Heat levels and associated risk
 - Exertion levels planned
 - o Acclimation level of those present for training
 - o <u>Time</u> factors including duration of activity and recovery time

Develop and implement risk controls

- Plan training in advance and estimate the risk of collapse events
- Consider the environment and training event characteristics:
 - Training event characteristics
 - Uniform and equipment needed
 - Location and time of day of key activities
- Adjust activity distances, durations, pace, and loads and emplace work/rest cycles
 - o Conduct high-intensity training in cooler morning hours
 - o Ensure no consecutive days of high-intensity activity
 - Ensure proper resources are at appropriate locations

- Equipment for checking Wet Globe Bulb Temperatures
- o Water, snacks and electrolyte beverages
- Medical resources and locations, and relevant communication capabilities
- Establish standard operating procedures, and train and test soldiers on them
- Identify how to monitor hydration & high risk personnel

Supervise and evaluate

- Ensure implantation of all planned measures
- Check junior leaders for knowledge of the prevention program
- Check medical resources (e.g., equipment, communications)
- Stop training if an injury occurs and re-assess the total unit status

Further clinician actions²

- Screen and defer individuals seeking entry to military service for histories of prior recurrent or severe exertional collapse
- Emplace protective duty restrictions for individuals with such histories
- Refer soldiers with such histories for review for possible discharge from service

References

- U.S. Army Medical Department, Army Public Health Center. Heat illness
 prevention. 2016.

 (https://phc.amedd.army.mil/topics/discond/hipss/pages/heatinjuryprevention.aspx)
- Department of the Army. Army Regulation 40-501: Standards of Medical Fitness.
 2011 (http://armypubs.army.mil/epubs/pdf/r40_501.pdf)